



MAX-PLANCK-GESELLSCHAFT

Wall adsorption of a colloidal particle moving in a quiescent partially wetting fluid




Alvaro Domínguez, Siegfried Dietrich

Max-Planck-Institut für Metallforschung & ITAP–University of Stuttgart, Germany

In thermal equilibrium, a colloidal particle between two parallel plates immersed in a fluid which partially wets both the particle and the plates, is attracted by the walls. However, if the particle moves parallel to the plates, a hydrodynamic lift force away from the plates arises in the limit of low Reynolds number. We study theoretically the competition of these two effects and identify the range of velocity in which the velocity may serve as a parameter controlling the adsorption in microflows.

Geometrical and physical setting



A spherical particle moves in a quiescent fluid with constant velocity parallel to two infinite plates

- Static interaction via dispersion force. Atomic potentials:

* Liquid-liquid, solid-solid: $\mathcal{V}_0(r) = -4\varepsilon(\sigma/r)^6$ ($\varepsilon \approx kT$)

* Liquid-solid: $\mathcal{V}_{ls}(r) = A\mathcal{V}_0(r)$

Relative Hamaker constant: $0 \leq A \leq 1$

A controls the wetting behavior. Contact angle: $\cos \theta \approx 2A - 1$

- Stationary incompressible flow:

$$\nabla \cdot \mathbf{v} = 0 \quad \nabla^2 \mathbf{v} - \frac{R}{\mu V} \nabla(p + W_{\text{liquid}}) = Re(\mathbf{v} \cdot \nabla) \mathbf{v}$$

and no-slip boundary conditions


Potential energy of the fluid in the dispersion field: W_{liquid}

Particle Reynolds number: $Re = \varrho R V / \mu \lesssim 1$

- Effective potential energy of the particle, $W(h) =$

hydrodynamic+dispersion =

$$kT \frac{R}{\sigma} Re^2 w_1\left(\frac{h}{R}, \frac{L}{R}, Re\right) - (1 - A) \varepsilon w_2\left(\frac{h}{R}, \frac{L}{R}\right)$$



- Motion in the h -direction: Brownian + relaxational dynamics:

$$6\pi\mu R v \lambda(h) + \frac{dW}{dh} = 0, \quad \lambda(h) = \text{wall-correction to Stokes drag}$$

- The Fokker-Planck equation for the particle motion yields the mean first-passage time, $\mathcal{T}$, from the midplane, $h = L/2 - R$, to wall contact, $h \approx \sigma$:

$$\mathcal{T} \propto \int_{L/2-R}^{\sigma} dy \int_{L/2-R}^y dz \lambda(y) \exp[W(y) - W(z)] / kT$$


- $\mathcal{T}$ is normalized to $\mathcal{T}_{\text{free}} =$ passage time in the absence of forcing:

$\mathcal{T} < \mathcal{T}_{\text{free}} \Rightarrow$ enhanced adsorption

$\mathcal{T} > \mathcal{T}_{\text{free}} \Rightarrow$ delayed adsorption

Results

- We study $\mathcal{T}$ as a function of A , Re and R for values of the parameters σ , ϱ , ε , μ appropriate for atomic liquids




- The qualitative shape of the function $\mathcal{T}(A, Re \lesssim 1)$ is independent of R , L in the range

$$10 \lesssim R/\sigma \lesssim 10^4 \quad 3 \lesssim L/R \lesssim 10$$

- The sensitivity of $\mathcal{T}$ to changes in Re increases with Re and R/σ

- Define a critical Re_c by the condition $\mathcal{T} = \mathcal{T}_{\text{free}}$:



- Some numbers: for atomic liquids $V(\text{m s}^{-1}) \sim Re/R(\mu\text{m})$

$$R \sim 1 \mu\text{m} \Rightarrow Re_c \sim 0.04 \Rightarrow V_c \sim 4 \text{ cm s}^{-1}$$

Conclusions

- In the **μ -meter scale**, the time of adsorption by the wall can be controlled in a realistic range of values of the particle velocity.
- In the **nano-meter regime**, the theory presented is pushed to its limits and suggests that the hydrodynamic lift is irrelevant.
- In particular, hydrodynamic lift is unlikely to explain the results of the MD simulation by Drazer et al., PRL **89** (2002) 244501.